

**Research Article**

# Pteridohytic flora of eastern edge of the Western Ghats in Kosergode, Kerala

**M. Maridass**

FISSD'S- Research Institute of Conservation Ecology (Run by Foundation for Innovative Science and Socio Economic Development (FISSD'S), Arunagiri Complex, Palayamkottai Tirunelveli, Tamilnadu-627002,India.

Email: drmaridass@gmail.com; Mobile : +91-9487567793

**ABSTRACT**

India boasts a rich diversity of pteridophyte flora, encompassing around 1200 species, a significant number of which are endemic. Key regions for pteridophyte diversity include the Himalayas, Eastern Ghats, and Western Ghats. These plants inhabit a range of environments, from sea level to mountainous areas, and are crucial to ecosystems while also possessing medicinal and economic significance. Pteridophytes constitute an essential part of the ecosystem, and since the majority of them reside in forests, they serve as effective indicators of issues such as deforestation and habitat loss. Numerous pteridophytes are included in the list of threatened species across various categories; however, effective solutions to address this global issue have yet to be explored. The objective of the current study was to investigate the pteridophytic diversity along the eastern edge of the Western Ghats in Kosergode, Kerala. The diversity assessment revealed that 87 species of pteridophytes were identified, belonging to 22 families as documented in this study.

**Keys:** Pteridophytes, diversity, Kasercode, Kerala, Western Ghats

**Maridass,M.(2025).** Pteridohytic flora of eastern edge of the Western Ghats in Kosergode, Kerala. *Pteridological Research*, Vol.No.14(1):1-7. (January-June)

**Article History**

Article received dated on : 29-11-2024

Corrected and Resubmission dated on 02– 12-2024

Published Online : 15-06-2025



## 1. INTRODUCTION

The term "Pteridophyta" is originated from Greek word "Pteron" which means feather and "phyton" means plants i.e. the plants having feather like fronds. The pteridophytes formed a dominant part of earth's vegetation about 282-230 million years ago during the Palaeozoic era. Late Palaeozoic era can be clearly regarded as the age of pteridophytes because during this period, pteridophytes were the dominant land plants on earth. There are about 13,000 species of pteridophytes across the world and 1,500 species of pteridophytes at random from a global checklist and carefully assessed the IUCN Red List status of each species. About 9% world Pteridophytes occurs in India or only in 2.5% landmass of the world. Western Ghats supports 349 pteridophytic species out of 1100-1200 species of ferns and fern allies in India (Manickam and Irudayaraj, 1992).

In India Pteridophytes are distributed in all the phytogeographical zones of India ranging from sea level to alpine Himalayas where they grow as Hydrophytes, Mesophytes, Lithophyte, Epiphyte, Hemiepiphyte, Climbers etc. They can be found in all ground habitats such as Ravine, Forest floor, on slopes, Grassland, on Rocks and crevices, on open walls and stone boulders and at certain places they form gregarious Thickets. As epiphytes different species of Pteridophytes also distributed on different part of tree as on base of tree, bole, branches, forking etc. Most of the pteridophytes diversity is observed in the Himalayas, Eastern and Western Ghats. Previous studies, worked on the pteridophytes of Western Ghats in detailed described in prominent publications are Ecological studies on the Fern Flora of Palni hills (Manickam, 1984), Fern Flora of Palni Hills, South India (Manickam, 1986), Cytology of the ferns of Western Ghats, South India (Manickam and Irudayaraj 1988) and Pteridophyte flora of Western Ghats, South India (Manickam and Irudayaraj, 1992).

Kasaragod district is situated on the eastern edge of the Western Ghats. The Western Ghats form an almost continuous mountain wall on the eastern side of the district, separating it from other areas outside of Kerala. Western Ghats in Kasaragod are home to various natural attractions, including the Kottancheri Hills, which are considered a trekker's paradise. The district also features the Ranipuram Wildlife Sanctuary, known for its diverse flora and fauna. The Mogral River originates in the lower foothills of the Western Ghats within Kasaragod. The Western Ghats in Kasaragod contribute to the district's rich biodiversity, with forests, hills, and rivers providing a habitat for various species of wildlife and plants. Aim of the present study was exploration of pteridophytic diversity of eastern edge of Western Ghats of Kosergode, Kerala from the period of 2022-2023.

## 2. Materials and Methods

### 2.1 Study area

Kasaragod district in Kerala is known for its close proximity to the Western Ghats, a major mountain range that runs along the Indian peninsula's western coast. The Western Ghats provide Kasaragod with a variety of natural attractions, including forests, hills, and rivers. The district itself is often referred to as the "Land of Gods"

and "Sapta Bhasha Sangama Bhumi" (Land of Seven Languages), highlighting its natural beauty and diverse cultural heritage. Extensive fieldwork was conducted in Kasaragod district in Kerala (February 2023). During the field survey of pteridophyte species, the specimen population, habitat, morphological characteristics, and types of forest/ponds were documented. The collected pteridophytes were identified using various literature photographs and specifications (Manickam and Irudayaraj, 1992).



Fig.1: Map of Kasaragod district of the Western Ghats.



**Table-1:** Pteridophytic diversity of Kasarcode regions of Kerala State.

Sl.No.	Species	Current Status	Family
1	<i>Asplenium adiantum-nigrum</i> L.	Common	Aspleniaceae
2	<i>Asplenium nidus</i> L.	Common	Aspleniaceae
3	<i>Asplenium longissimum</i> Blume	Common	Aspleniaceae
4	<i>Asplenium scolopendrium</i> L.	Common	Aspleniaceae
5	<i>Athyrium filix-femina</i> (L.) Roth	rare	Athyriaceae
6	<i>Athyrium hohenackerianum</i> (Kze.) Moore	Common	Athyriaceae
7	<i>Athyrium asplenoides</i> (Michx.) Desv.,		Athyriaceae
8	<i>Blechnum orientale</i> L.	rare	Blechnaceae
9	<i>Blechnum cartilagineum</i> Sw.	rare	Blechnaceae
10	<i>Cyathea nilgirensis</i> Holttum	endemic	Cyatheaceae
11	<i>Cyathea spinulosa</i> Wall ex Hook	endemic	Cyatheaceae
12	<i>Bolbitis semicordata</i> (Bak.) Ching	endemic	Dryopteridaceae
13	<i>Arachniodes sledgei</i> Fraser-Jenk.	Common	Dryopteridaceae
14	<i>Diplazium acrostichoides</i> (Sw.) Butters	Common	Dryopteridaceae
15	<i>Rumohra adiantiformis</i> (G. Forst.) Ching	Common	Dryopteridaceae
16	<i>Cystopteris fragilis</i> (L.) Bernh.	Common	Dryopteridaceae
17	<i>Pteridium revolutum</i> (Blume) Nakai	Common	Dennstaedtiaceae
18	<i>Microlepia firma</i> Mett. ex Kuhn	Common	Dennstaedtiaceae
19	<i>Microlepia speluncae</i> (L.) T. Moore	Common	Dennstaedtiaceae
20	<i>Pteridium aquilinum</i> subsp. <i>wightianum</i> (Wall. ex J. Agardh) ...	Common	Dennstaedtiaceae
21	<i>Gleichenia dicarpa</i> R.Br.	Common	Gleicheniaceae
22	<i>Dicranopteris linearis</i> (Burm. f.) Underw		Gleicheniaceae
23	<i>Trichomanes companulatum</i> Roxb	Common	Hymenophyllaceae
24	<i>Hymenophyllum denticulatum</i> Sw.	Common	Hymenophyllaceae
25	<i>Trichomanes christii</i> Copel	Common	Hymenophyllaceae
26	<i>Trichomanes lunulatum</i> (Madhus. & C.A.Hameed)	Endemic	Hymenophyllaceae
27	<i>Trichomanes saxifragoides</i> C.Presl	Common	Hymenophyllaceae
28	<i>Trichomanes christii</i> Copel	Common	Hymenophyllaceae
29	<i>Isoetes coromandeliana</i> L. f	Common	Isoetaceae
30	<i>Lindsaea malabarica</i> (Bedd.) Baker ex Christ	endemic	Lindsaeaceae
31	<i>Lindsaea ensifolia</i> Sw.	Common	Lindsaeaceae
32	<i>Lygodium microphyllum</i> (Cav.) R.Br.	Common	Lygodiaceae
33	<i>Lygodium flexuosum</i> (L.) Sw.	common	Lygodiaceae
34	<i>Marsilea quadrifolia</i> L.	Common	Marsileaceae
35	<i>Marsilea minuta</i> L.	Common	Marsileaceae
36	<i>Psilotum nudum</i> (L.) P.Beauv.	endemic	Psilotaceae
37	<i>Pyrrosia porosa</i> var <i>Porosa</i> Hovenkamp	Common	Polypodiaceae



38	<i>Pyrrosia lanceolata</i> (L.) Farw. <i>Synonyms:</i> <i>Acrostichum lanceolatum</i> L.	Common	Polypodiaceae
39	<i>Pyrrosia heterophylla</i> (L.) M. G. Price	Common	Polypodiaceae
40	<i>Drynaria quercifolia</i> Linn. J. smith	Common	Polypodiaceae
41	<i>Tectaria zeilanica</i> (Houtt.) Sledge	endemic	Tectariaceae
42	<i>Tectaria coadunata</i> (Wall. ex Haines) Raizada & N.P. ...	Common	Tectariaceae
43	<i>Thelypteris caudipinna</i> Ching	Common	Thelypteridaceae
44	<i>Macrothelypteris torresiana</i> (Gaudich.) Ching	Common	Thelypteridaceae
45	<i>Thelypteris dentata</i> (Forssk.) E.P.St.John (Syn.) <i>Christella dentata</i> (Forssk.) Brownsey & Jermy	Common	Thelypteridaceae
46	<i>Thelypteris trigonospora</i> (Holtum) Fraser-Jenk.S	Common	Thelypteridaceae
47	<i>Aleuritopteris rufa</i> (D.Don) Ching - (Syn.) <i>Cheilanthes rufa</i> (D.Don)	endemic	Pteridaceae
48	<i>Adiantum latifolium</i> Lam.	Common	Pteridaceae
49	<i>Adiantum incisum</i> C.Presl.	Common	Pteridaceae
50	<i>Adiantum tenerum</i> Sw.	Common	Pteridaceae
51	<i>Adiantum philippense</i> L.	Common	Pteridaceae
52	<i>Adiantum caudatum</i> L.	Common	Pteridaceae
53	<i>Adiantum abscissum</i> Schrad.	Common	Pteridaceae
54	<i>Adiantum capillus-veneris</i> L.	Common	Pteridaceae
55	<i>Acrostichum aureum</i> L.	Common	Pteridaceae
56	<i>Adiantum lunulatum</i> Burm.fil.	Common	Pteridaceae
57	<i>Adiantum raddianum</i> Presl.	rare	Pteridaceae
58	<i>Parahemionitis cordata</i> (Roxb. ex Hook. & Grev.) Fraser-Jenk.	common	Pteridaceae
59	<i>Pityrogramma calomelanos</i> (L.) Link	Common	Pteridaceae
60	<i>Pteris argyrea</i>	Common	Pteridaceae
61	<i>Pteris biaurita</i> L.	Common	Pteridaceae
62	<i>Peris pellucida</i> C.Presl	Common	Pteridaceae
63	<i>Pteris vittata</i> L.	Common	Pteridaceae
64	<i>Pteris ensiformis</i> Burm.f.	Common	Pteridaceae
65	<i>Pteris chilensis</i> Desv.	Common	Pteridaceae
66	<i>Ceratopteris thalictroides</i> (L.) Brongn.	Common	Pteridaceae
67	<i>Cheilanthes tenuifolia</i> (Burm. f.) Sw.	rare	Pteridaceae
68	<i>Actiniopteris radiata</i> (Koenig ex Sw.) Link	Common	Pteridaceae
69	<i>Salvinia herzogii</i>	Common	Salviniaceae
70	<i>Salvinia minima</i> Baker	Common	Salviniaceae
71	<i>Salvinia natans</i> (L.) All.	Common	Salviniaceae
72	<i>Salvinia rotundifolia</i> Wild.	Common	Salviniaceae
73	<i>Azolla pinnata</i> R.Br.	Common	Salviniaceae
74	<i>Nephrolepis exaltata</i> (L.) Schott	Common	Nephrolepidaceae
75	<i>Nephrolepis auriculata</i> (L.)	Common	Nephrolepidaceae
76	<i>Nephrolepis cordifolia</i> (L.) C.Presl	Common	Nephrolepidaceae



77	Osmunda huegeliana C.Presl	rare	Osmundaceae
78	Osmunda regalis L.	Common	Osmundaceae
79	Ophioglossum reticulatum L.	rare	Ophioglossaceae
80	Ophioglossum madhusoodanum Sojan	rare	Ophioglossaceae
81	<i>Ophioglossum raphaelianum</i> Anto P.V.	Common	Ophioglossaceae
82	<i>Selaginella ciliaris</i> (Retz.) Spring	rare	Selaginellaceae
83	<i>Selaginella involvens</i> (Sw.) Spring	rare	Selaginellaceae
84	<i>Selaginella kraussiana</i> (Kunze) A.Braun	rare	Selaginellaceae
85	<i>Selaginella delicatula</i> (Desv.) Alston.	Common	Selaginellaceae
86	<i>Selaginella wildenovii</i> (Desv. ex Poir.) Baker	Common	Selaginellaceae
87	<i>Selaginella ciliaris</i> (Retz.) Spring	Common	Selaginellaceae

Table-2: Total number of Families and species of forest regions of Kasargode District, Kerala

Sl.No.	Family	No. of Species
1	Aspleniaceae	4
2	Athyriaceae	3
3	Blechnaceae	2
4	Cyatheaceae	2
5	Dryopteridaceae	5
6	Dennstaedtiaceae	4
7	Gleicheniaceae	2
8	Hymenophyllaceae	6
9	Isoetaceae	1
10	Lindsaeaceae	2
11	Lygodiaceae	2
12	Marsileaceae	2
13	Psilotaceae	1
14	Polypodiaceae	4
15	Tectariaceae	2
16	Thelypteridaceae	4
17	Pteridaceae	22
18	Salviniaceae	5
19	Nephrolepidaceae	3
20	Osmundaceae	2
21	Ophioglossaceae	3
22	Selaginellaceae	6



### 3. Results and Discussion

The current field study is a survey of Pteridophytes located at the eastern edge of the Western Ghats Forest region in Kasargode, conducted during the period from June 2022 to February 2023. The findings of this study indicate that 87 species were identified, belonging to 22 families, as recorded from the regions of Kasargode (see Table 1 and 2). The most of the Pteridophytes species were observed by terrestrial, epiphytic and lithophytic forms found this forest. I have observed the present study revealed that three endemic species identified such as *Cyathea nilgiriensis*, *Cyathea spinulosa* Wall ex Hook and *Bolbitis semicordata*. Previous studies have included several reports of research articles focusing on the pteridophytic diversity in the eastern and western Ghats regions. Holtum (1976) documented 10 members of Thelypteridaceae in the "Flora of Hassan District." Yoganarasimhan et al. (1981) identified 12 species of ferns in their "Flora of Chikmagalur District" in Karnataka. According to Blatter and Almeida (1992), 90 species of ferns and fern allies were reported from the Uttara Kannada district. Ramachandra et al. (2010) recorded 54 pteridophyte species from the Gundia river basin in Hassan district. Previous research conducted by Subina et al. (2021) indicated that 16 families across 24 genera and 44 species were documented in Pampadum Shola National Park. In a separate study, Abhilaksha (2023) noted that a total of 152 species of pteridophytes, classified into 62 genera and 26 families, were identified across various states including Uttar Pradesh, Uttarakhand, Himachal Pradesh, and Jammu and Kashmir.

The current research identifies the predominant families of Pteridaceae (22 species), Hymenophyllaceae (6 species), and Selaginellaceae (6 species) within the forest regions of Kasargode (Fig.1). According to Dixit (2000), an analysis of species diversity at the family level reveals that the highest diversity is found in the Aspleniaceae family, which includes 27 species, followed by Polypodiaceae (25 species), Athyriaceae (24 species), Thelypteridaceae (23 species), Selaginellaceae (20 species), Pteridaceae (17 species), Aspidaceae (13 species), among others. At the generic level, the greatest diversity is noted in the genera Selaginella (20 species), Pteris (17 species), and Diplazium (7 species). This study has confirmed that the dominant families are Pteridaceae, Hymenophyllaceae, and Selaginellaceae, along with the presence of rare and endemic species of pteridophytes. The lady fern genus *Athyrium* represents one of the most diversified lineages in Athyriaceae with about 160-220 known species (Ran Wei et al., 2018). I have identified three species of *Athyrium* sp in Kasargode forest in Kerala. Praveen Kumar and Udayan (2018) reported that *Salvinia molesta* and *Marsilea minuta* as their wide distribution as weeds in ponds and paddy fields Kerala forest regions. According to Iwatsuki (1990) reported that Hymenophyllaceae comprises more than 600 species of delicate ferns that are most abundant in humid tropical forests, but extend in humid shady habitats into temperate regions. I have detailed investigation of present study 6 species of Hymenophyllaceae family such as *Trichomanes complanatum* Roxb; *Hymenophyllum denticulatum* Sw.; *Trichomanes christii* Copel; *Trichomanes lunulatum* (Madhus. & C.A.Hameed); *Trichomanes saxifragoides* C.Presl; and *Trichomanes christii* Copel.

According to Dixit, (2000) reported that due to unplanned felling of trees in the forests the members of epiphytic pteridophytes belonging to the families Polypodiaceae, Davalliaeae, Aspleniaceae and Vittariaceae, have been reduced. Earlier studies, Sumesh Dudani et al., (2002) reported that serious threat to the several pteridophytes species like *Psilotum nudum*, *Tectaria zeylanica*, *Lindsaea malabarica*, and *Cheilanthes rufa*.

### Conclusion

Conclusion of the present study was many rare and endangered ferns and fern-allies like *Psilotum nudum*, *Tectaria zeylanica*, *Lindsaea malabarica*, *Cheilanthes rufa*, and *Cyathea nilgiriensis*, etc. have been recorded from the Western Ghats and these species were need to urgent attention for conservation.

### 4. Conflicts of Interest

The author declare that they have no conflicts of interest.

### 5. References

Bunyan, M., Bardhan, S., Singh, A. & Jose, S. (2015). Effect of topography on the Tropical montane forest fragments: a predictive modelling approach. *Journal of Tropical Forest Science*, 27(1): 30-38.

Swarupanandan, K., Sasidharan, N., Chacko, K.C. & Chand, B.S. (1998). Studies on the Shola forests of Kerala. KFRI research report. Zar J.H (1999) Biostatistical analysis, 4th edn. Prentice Hall International, USA.

Blatter, E. and Almeida, J.F.D (1992). The Ferns of Bombay. D.B. Taporevala Sons & Co., Bombay.

Yoganarasimhan, S.N., Subramanyam, K. and Razi, B.A. (1981). Flora of Chikmagalur district. International Book distributors, Dehradun.

Rajagopal, P.K. and Bhat, K.G. (1998). Pteridophyte flora of Karnataka state, India. *Indian Fern Journal*, 15: 1-28.

Ramachandra, T.V., Chandran, M.D.S., Bhat, H.R., Dudani, S., Rao, G.R., Boominathan, M., Mukri, V. and Bharath, S. (2010). Biodiversity, Ecology and Socio-economic aspects of Gundia river basin in the context of proposed mega hydroelectric power project. CES Technical Report-122, Centre for Ecological Sciences, Indian Institute of Science, Bangalore.

Dixit, R.D. (2000). Conspectus of Pteridophytic diversity in India. *Indian Fern Journal*, 17:77 - 91.

Manickam, V.S. (1984). Ecological studies on the fern flora of the Palni hills (South India). Today and Tomorrow's Printers and Publishers, New Delhi.

Manickam, V.S. and Irudayaraj, V. (1988). Cytology of the ferns of Western Ghats (South India). Today and Tomorrow's Printers and Publishers, New Delhi.

Manickam, V.S. and Irudayaraj, V. (1992). Pteridophyte Flora of the Western Ghats - South India. B. I. Publications, New Delhi.

Manickam, V.S. (1986). Fern flora of the Palni Hills (South India). Today and Tomorrow's Printers and Publishers, New Delhi.

Schneller, J.J. (1980). Cytotaxonomic investigations of *Salvinia herzogii* de la Sota. *Aquatic Botany*, 9:279-283.

Abhilaksha,(2023). Exploring the Rich Biodiversity of Pteridophytes in Northern India: A Study. *Int. J. Sci. Res. Sci & Technol.*, 10 (2): 567-594.



Manickam, V.S. and Irudayaraj, V. (1992). Pteridophyte Flora of the Western Ghats- South India. B. I. Publications, New Delhi.

Sumesh Dudani, Subhash Chandran, M.D. and Ramachandra, T.V. (2002). Pteridophytes of Western Ghats. Biodiversity Documentation and Taxonomy, Pages 343-351. Edited by: A. Biju Kumar, Narendra Publishing House.

Ran Wei, Atsushi Ebihara, Yan-Mei Zhu, Cun-Feng Zhao, Sabine Hennequin, Xian-Chun Zhang, (2018). A total-evidence phylogeny of the lady fern genus *Athyrium* Roth (Athyriaceae) with a new infrageneric classification, Molecular Phylogenetics and Evolution, 119:25-36.

Praveen Kumar, K. and Udayan, P.S. (2018). A Survey on the Pteridophyte Flora of the 18 Selected Sacred Groves in Chalavara Grama Panchayath, Palakkad District, Kerala. International Journal of Environment, Agriculture and Biotechnology (IJEAB),3(4):1222-1225.

In this article designed and published © 2024 by GTRP  
 BIOJOURNALS is licensed under Creative Commons Attribution-Non Commercial - No Derivatives 4.0 International

